

Electronic Supplementary Material

Ongoing invasions of the African clawed frog, *Xenopus laevis* - a global review

G. J. Measey^{1*}, D. Rödder², S. L. Green³, R. Kobayashi⁴, F. Lillo⁵, G. Lobos⁶, R. Rebelo⁷ & J.-M. Thirion⁸

¹School of Environmental Sciences and Development, North-West University, Private Bag X6001, Potchefstroom 2520, South Africa. e-mail: john@measey.com

²Zoologisches Forschungsmuseum Alexander Koenig (ZFMK), Section of Herpetology, Adenauerallee 160, D-53113 Bonn, Germany. e-mail:

d.roedder.zfmk@uni-bonn.de

³Department of Comparative Medicine, School of Medicine, Stanford University, Stanford, California, USA. e-mail: sherril@stanford.edu

⁴Center for Toki and Ecological Restoration (CTER), Niigata University, 1101-1 Niibo-katagami, Sado-City, Niigata, 952-0103, Japan e-mail: raita@cc.niigata-u.ac.jp

⁵Dipartimento di Biologia Ambientale e Biodiveristà, Università` di Palermo, Via Archirafi 18, 90123 Palermo, Italy e-mail: francesco.lillo@gmail.com


⁶Centro de Vida Silvestre, Facultad de Ciencias Veterinarias y Pecuarias, Universidad de Chile, Casilla 2, Correo 15, La Granja, Santiago, Chile e-mail: galobos@ug.uchile.cl

⁷Centro de Biología Ambiental and Departamento de Biología Animal, Faculdade de Ciencias da Universidade de Lisboa, Campo Grande, 1749-016 Lisbon, Portugal e-mail: rmrebelo@fc.ul.pt

⁸Association Objectifs BIOdiversitéS (OBIOS), 22 rue du docteur Gilbert; 17250 - Pont-l'abbé-d'Arnoult, France. e-mail: thirion.jean-marc@sfr.fr


*Corresponding author: tel: +27 21 799 8860 fax: +27 21 797 6570
email: john@measey.com

Predicted distribution of *Xenopus laevis* with a spatial resolution of 2.5 arcmin from a Maxent model (see article text for details). Warmer colours indicate higher climatic suitability. Suitable climatic areas above the minimum training presence logistic threshold are indicated in light downward diagonal fill, wherein those optimal areas above the 10 % training omission threshold are indicated in dark downward diagonal fill. Areas with bioclimatic conditions exceeding those in the training areas requiring model extrapolation are indicated as crosshatched.




Legend

●	invasive	Maxent score	Thresholds
●	native	High : 0.92	> min train presence
▨	Extrapolation (MESS)	Low : 0.00	> 10% train omission




Legend

●	invasive	Maxent score	Thresholds
●	native	High : 0.92	< min train presence
▨	Extrapolation (MESS)	Low : 0.00	> 10% train omission



Legend

●	invasive	Maxent score	Thresholds
●	native	High : 0.92	< min train presence
▨	Extrapolation (MESS)	Low : 0.00	> 10% train omission



Legend

● invasive	Maxent score	Thresholds
● native	High : 0.92	< min train presence
▨ Extrapolation (MESS)	Low : 0.00	> 10% train omission